

Sample HelpDesk Answer Submission: **Diagnostic Question**

Manuscripts are submitted in the following format. To see how this manuscript appeared in print, please reference HDA Document #125 *Sample HDA Published: Diagnostic Question*.

2015 EBP

Title: Patient with leg cramps

Word Count: 487

What is the diagnostic approach to a patient with leg cramps?

Evidence-Based Answer

A careful history targeting possible peripheral vascular disease, arthritis, and medication use (particularly diuretics, statins, and long-acting bronchodilators) is likely helpful in evaluating a patient with leg cramps. (SOR: **B**, observational studies). Laboratory evaluation is not known to be helpful (SOR: **C**, extrapolated from studies in runners).

Evidence Summary

A 1999 cross-sectional study of adults 65 years of age and older (N=365) investigated the prevalence of leg cramps and their association with specific disease conditions.¹ Data were collected from patient questionnaires administered over a 6-month period. The overall prevalence of leg cramps was 50%. Leg cramps were most prevalent at night (62%). Patients with leg cramps showed a strong association with peripheral vascular disease (OR 2.9; 95% CI, 1.9–4.6), arthritis (OR 2.3; 95% CI, 1.5–3.5), and female sex (OR 2.0; 95% CI, 1.3–3.0) compared with individuals without leg cramps.¹

A 2012 sequence symmetry analysis sought to determine if treatment for muscle cramps increased during the year after initiation of statin, diuretic, or long-acting β_2 -agonist (LABA) therapy.² By searching healthcare databases in British Columbia and Canada, the number of new quinine prescriptions written (initiated for treatment of cramps) were identified before and after the initiation of treatment with the index medications, in adults 50 years or older. Adjusted sequence ratios (ASR) showed initial quinine prescriptions were significantly more likely to follow rather than precede new prescriptions for all index medications: diuretics (ASR 1.5; 95% CI, 1.3–1.6), statins (ASR 1.2; 95% CI, 1.0–1.3), and LABAs (ASR 2.4; 95% CI, 2.0–2.9). Both thiazide-like diuretics (ASR 1.5; 95% CI, 1.3–1.7) and loop diuretics (ASR 1.2; 95% CI, 1.0–1.4) were associated with increased quinine prescriptions. Thus in patients presenting with complaints of leg cramps, the possibility of diuretics, statins, and LABAs as etiologic agents should be explored.²

A 2003 individual cohort trial evaluated the development of exercise-associated muscle cramping (EAMC) in 72 long-distance runners.³ The runners were divided into those with and without a history of EAMC. Lab studies (Na, K, Ca, Mg, and hemoglobin) and weight were

measured pre-race, immediately postrace (56 km), and 60 minutes post-race. The only laboratory measurements showing statistically significant differences between the groups were a lower immediate post-race serum sodium concentration in the cramp group versus the control group (mean 140 vs 142 mEq/L; P=.004) and higher immediate post-race magnesium level (1.7 vs 1.6 mg/dL; P=.03).

Matthew Nix, MD

Zoha Iqbal, MD

Brian Caldwell, MD, MS

Michael Downs, MD

U of AR for Medical Sciences-AHEC, Southwest
Texarkana, AR

1. Abdulla AJ, Jones PW, Pearce VR. Leg cramps in the elderly: prevalence, drug and disease associations. *Int J Clin Pract.* 1999; 53(7):494–496. **[STEP 3]**
2. Garrison SR, Dormuth CR, Morrow RL, et al. Nocturnal leg cramps and prescription use that precedes them: a sequence symmetry analysis. *Arch Intern Med.* 2012; 172(2):120–126. **[STEP 3]**
3. Schwellnus MP, Nicol J, Laubscher R, et al. Serum electrolyte concentrations and hydration status are not associated with exercise associated muscle cramping (EAMC) in distance runners. *Br J Sports Med.* 2004; 38(4):488–492. **[STEP 2]**