



Keri Bergeson, MD;
Shailendra Prasad, MBBS,
MPH; Janice Benson, MD
North Memorial
Family Medicine Residency,
University of Minnesota,
Minneapolis (Drs. Bergeson
and Prasad); University
of Chicago (NorthShore)
Department of Family
Medicine (Dr. Benson)

PURLS EDITOR

Anne Mounsey, MD
Department of Family
Medicine, University of
North Carolina at Chapel
Hill

Skip the compression stockings following DVT

Although commonly used, compression stockings do not effectively prevent post-thrombotic syndrome.

PRACTICE CHANGER

Do not recommend elastic compression stockings (ECS) to decrease the incidence of post-thrombotic syndrome (PTS) after deep vein thrombosis (DVT).¹

STRENGTH OF RECOMMENDATION

B: Based on a large, randomized controlled trial

Kahn SB, Shapiro S, Wells PS, et al; SOX trial investigators. Compression stockings to prevent post-thrombotic syndrome: a randomised placebo-controlled trial. *Lancet*. 2014;383:880-888.

ILLUSTRATIVE CASE

A 56-year-old man comes to your clinic 3 days after receiving a diagnosis of lower extremity deep vein thrombosis (DVT). He was prescribed warfarin, 5 mg/d, with enoxaparin bridging, 120 mg/d. He has read about post-thrombotic syndrome (PTS) online and is very concerned about this possible side effect. He is asking about using elastic compression stockings (ECS). What should you tell him?

PTS can be a frustrating, debilitating condition. Its clinical features range from minor limb swelling to severe edema and pain, irreversible skin changes, and leg ulcerations.² It occurs in 25% to 50% of patients after DVT.³ Because current PTS treatments are not very effective, prevention is essential.^{4,5}

Patients are frequently encouraged to wear ECS after DVT to reduce the incidence of PTS by reducing venous hypertension and reflux. These stockings are expensive and uncomfortable. Prior studies suggested

that using ECS can cut the incidence of PTS in half.^{6,7} However, these were small, single-center studies, and they were not placebo-controlled.^{6,7}

STUDY SUMMARY

RCT sets aside a common practice

Kahn et al¹ conducted a randomized, placebo-controlled trial of active vs placebo ECS in patients from 24 centers in the United States and Canada who'd had an ultrasound-confirmed proximal DVT (in the popliteal or more proximal deep leg vein) within the previous 14 days. Most patients received standard anticoagulation therapy to treat their DVT (5-10 days of heparin and 3-6 months of warfarin). Patients were excluded if they had received thrombolytics, had arterial claudication, had a life expectancy of <6 months, were unable to put on ECS due to physical disabilities or allergy, or were unable to participate in follow-up visits.

Patients were randomly assigned to wear active (30-40 mm Hg graduated) ECS or identical-looking placebo ECS with <5 mm Hg compression at the ankle for 2 years. Providers, study personnel and statisticians, and patients were all blinded to treatment allocation. Patients were asked to wear the stocking on the affected leg each day from when they woke until they went to bed.

Participants were followed at one, 6, 12, 18, and 24 months. The primary outcome was the cumulative incidence of PTS diagnosed at 6 months or later using Ginsberg's



INSTANT POLL

How often do you recommend compression stockings for patients who have developed deep vein thrombosis?

- Always
- Only for patients at high risk for post-thrombotic syndrome
- Only if patients ask about them
- Never

jfponline.com

criteria of ipsilateral pain and swelling of at least 1 month's duration.⁸ Secondary outcomes included severity of PTS, presence of leg ulcers, recurrence of venous thromboembolism (VTE), death, adverse events, venous valvular reflux, and quality of life (QOL). Outcomes were measured objectively using a validated scale (the Villalta scale) for PTS severity and the 36-item Short Form Health Survey (SF-36) and the Venous Insufficiency Epidemiological and Economic Study Quality of Life (VEINES-QOL) questionnaire to measure QOL.⁹⁻¹¹

There were 409 patients in the ECS group and 394 in the placebo group. Baseline characteristics, including body mass index (BMI), VTE risk factors, and anticoagulation treatment regimens, were similar between groups. The average age was 55.4 years in the study group (standard deviation [SD] ± 15.3 years) and 54.8 years (SD ± 15.8 years) in the placebo group. Men comprised 62.4% of the active group and 57.9% of the placebo group. Approximately 90% of the participants in both groups were white.

At one month, approximately 95% of participants in both the active and placebo groups used the stockings; at 24 months, a little less than 70% of the participants in both groups continued to use the stockings. The percentage of people who used the stockings for at least 3 days a week was similar across both groups.

The cumulative incidence of PTS during follow-up was 14.2% in the active group vs 12.7% in the placebo group, with a hazard ratio of 1.13 (95% confidence interval [CI], .73-1.76; *P*=.58). There were no differences in any of the secondary outcomes. Prespecified subgroup analyses found that age, BMI, and severity of DVT had no effect on the outcomes. There was a marginal benefit for ECS for women (*P*=.047) over men, but this does not likely reflect a true difference because the CIs surrounding the hazard ratios for men and women overlapped and crossed the null value.

WHAT'S NEW

New evidence contradicts previous studies

Two prior studies showed that using 30 to

40 mm Hg ECS decreased the incidence of PTS after proximal DVT.^{6,7} However, these were smaller, open-label, single-center studies. This study by Kahn et al¹ was the first placebo-controlled, randomized, multicenter study that used validated instruments to measure PTS and QOL. It found no benefit in using ECS, thus contradicting the results of the prior studies.

There are currently no guidelines or consensus statements for or against the use of ECS after DVT.

CAVEATS

High nonadherence rates might have affected the results

In both groups, adherence to the assigned intervention diminished throughout the study. Overall, approximately 95% of patients reported wearing their stockings at one month; this dropped to just under 70% by 2 years. Theoretically, this could have affected efficacy outcomes. However, the decrease was similar in both groups and represents what is observed in clinical practice. A prespecified per protocol analysis of patients who wore their ECS more regularly found no benefit.

It is possible that a "placebo effect" could explain the lack of difference between groups. However, the placebo stockings provided virtually no compression, and the 2-year cumulative incidence of PTS in both the treatment and placebo groups was similar to that seen in control groups in prior studies.^{6,7}

Finally, the incidence of PTS in this study was much lower than the 25% to 50% incidence reported in previous studies. Kahn et al¹ suggested that this was because they used more stringent and standardized criteria for PTS than was used in previous research.

CHALLENGES TO IMPLEMENTATION

There are no barriers to ending this practice

We see no challenges to implementation of this recommendation. **JFP**

ACKNOWLEDGEMENT

The PURLs Surveillance System was supported in part by Grant



This placebo-controlled randomized trial found no benefit in using compression stockings to prevent post-thrombotic syndrome.

Number UL1RR024999 from the National Center For Research Resources, a Clinical Translational Science Award to the University of Chicago. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center For Research Resources or the National Institutes of Health.

Copyright © 2014. The Family Physicians Inquiries Network. All rights reserved.

References

1. Kahn SR, Shapiro S, Wells PS, et al; SOX trial investigators. Compression stockings to prevent post-thrombotic syndrome: a randomized placebo-controlled trial. *Lancet*. 2014;383:880-888.
2. Kahn SR, Shrier I, Julian JA, et al. Determinants and time course of the postthrombotic syndrome after acute deep venous thrombosis. *Ann Intern Med*. 2008;149:698-707.
3. Prandoni P, Lensing AW, Cogo A, et al. The long-term clinical course of acute deep venous thrombosis. *Ann Intern Med*. 1996;125:1-7.
4. Cohen JM, Akl EA, Kahn SR. Pharmacologic and compression therapies for postthrombotic syndrome: a systematic review of randomized controlled trials. *Chest*. 2012;141:308-320.
5. Henke PK, Comerota AJ. An update on etiology, prevention, and therapy of postthrombotic syndrome. *J Vasc Surg*. 2011;53:500-509.
6. Brandjes DP, Büller HR, Heijboer H, et al. Randomised trial of effect of compression stockings in patients with symptomatic proximal-vein thrombosis. *Lancet*. 1997;349:759-762.
7. Prandoni P, Lensing AW, Prins MH, et al. Below-knee elastic compression stockings to prevent the post-thrombotic syndrome: a randomized, controlled trial. *Ann Intern Med*. 2004;141:249-256.
8. Ginsberg JS, Hirsh J, Julian J, et al. Prevention and treatment of postphlebotic syndrome: results of a 3-part study. *Arch Intern Med*. 2001;161:2105-2109.
9. Villalta S, Bagatella P, Piccioli A, et al. Assessment of validity and reproducibility of a clinical scale for the post-thrombotic syndrome. *Haemostasis*. 1994;24:158a.
10. McHorney CA, Ware JE Jr, Raczek AE. The MOS 36-Item Short-Form Health Survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs. *Med Care*. 1993;31:247-263.
11. Kahn SR, Lamping DL, Ducruet T, et al; VETO Study Investigators. VEINES-QOL/Sym questionnaire was a reliable and valid disease-specific quality of life measure for deep venous thrombosis. *J Clin Epidemiol*. 2006;59:1049-1056.

WE WANT TO HEAR FROM YOU!

Have a comment on an article, editorial, or department? You can send it by:

1. **E-MAIL:** jfp.eic@gmail.com
2. **FAX:** 973-206-9251 or
3. **MAIL:** The Journal of Family Practice, 7 Century Drive, Suite 302, Parsippany, NJ 07054

THE JOURNAL OF
FAMILY PRACTICE

LETTERS SHOULD BE 200 WORDS OR LESS. THEY WILL BE EDITED PRIOR TO PUBLICATION.

 Visit us @ jfponline.com

Watch & Learn:
A how-to video series
Cryosurgery



 **INSTANT POLL**

How often do you recommend compression stockings for patients who have developed deep vein thrombosis?

ONLINE EXCLUSIVES

- Pica: An age-old eating disorder that's often missed

PHOTO ROUNDS FRIDAY

Test your diagnostic skills at www.jfponline.com/articles/photo-rounds-friday.html



PLUS

Today's headlines in family medicine

GET UPDATES FROM US ON

FACEBOOK  **AND** **TWITTER** 

www.facebook.com/JFamPract <http://twitter.com/JFamPract>

www.jfponline.com