

Using CT effectively + flexibly: Conceptualize, motivate and intervene

By Dr. Claudia J. Haferkamp,
Graduate and Clinical Psychology Coordinator,
Millersville University

Workshop Overview

- **Hour 1.5: Core CBT elements + conceptualization:**
 - Maintaining reasonable structure + focus
 - Case conceptualization
- **Hour #1.5-3: Facilitating motivation + treatment integration + flexibility**
 - CBT + Motivational Interviewing
 - Exposure treatment: improving outcomes
 - Other issues? (time permitting)

CBT Core Elements

- Collaborative empiricism (Beck, 2011)
- Problem-focused
- Reasonably structured sessions
- Psycho-educational and...
- Focused on relapse prevention
- Action plans (homework) are essential

“Typical” CBT session structure

- Check in + mood check (+ objective data?)
- Collaboratively set initial session agenda
- Review action plan (homework)
- Re-prioritize final session agenda
- Discuss problems + **make interventions**
- Final session summary
- Review/develop next action plan (homework)
- Elicit CL feedback

Loss of session structure + focus

- Not adequately socializing CL to CBT
- Not discussing specific situations OR unfocused discussions:
 - Not focused on KEY thoughts, feelings, etc.
 - Unclear purpose of discussion OR no interventions made
 - TH's thoughts about interrupting
- Not eliciting or responding to CL feedback

Cbtscience/training/resources: Session assignment + feedback

- **Sample items: How well did:**
 - You feel heard + understood in today's session?
 - Today's session help address your problems?
 - How confusing was today's session?
 - How confident are you that you are progressing towards your therapy goals?

CBT Conceptualization

- **Can't treat what you can't conceptualize**
- Conceptualizations help us:
 - Organize CL info
 - Develop working hypotheses
 - Develop treatment plans + rationales for interventions
 - Build the working alliance

Why form working hypotheses? (Persons, 2015)

- One causal mechanism may underlie multiple problems
- ESTs may target a single disorder **ONLY**
 - There are no ESTs for many disorders
- Helps us address therapy-interfering thoughts + behaviors

Persons: Case Formulation

- Inclusive problem list
- Origins
- Precipitants (large events trigger current episode)
- Antecedents (activating situations): triggers symptoms
- Behaviors
- Consequences (functional)
- **Causal mechanisms (mostly cognitive)**
- Organismic variables (unique vulnerabilities)

Case formulation example (adapted) (Ledley et al., 2010)

- Mike's parents held him to exacting standards and sent him to schools known for academic rigor (**ORIGINS**). As a result, Mike started seeing others as critical and feared being rejected for making mistakes (**CAUSAL MECHANISMS**). These thoughts occurred after deciding to enter the priesthood (**PRECIPITANT**). Afterwards he had ATs such as, "I make more mistakes than others" and "people will notice my anxiety...they'll think I'm incompetent" resulting in increased (social) anxiety (i.e., blushing, sweating, sleep disruption) (**SYMPTOMS/PROBLEMS**).

Case formulation (cont.)

- Having to give his first sermon triggered his anxiety again (**ACTIVATING SITUATION**). Mike coped by over-preparing sermons, only spoke to familiar people at social events and avoided discussing the priesthood with his family (**MECHANISMS**). This temporarily reduced his anxiety (**FUNCTIONAL CONSEQUENCES**) but he missed making valuable social contacts and did not self-disclose with his mentors which left him feeling more dejected and unsure of his future (**SYMPTOMS/PROBLEMS**).

Inclusive problem lists


- **Problem areas:**
 - Health, psychiatric
 - Interpersonal/family
 - Job/school
 - Financial
 - Housing
 - Legal
 - Leisure functioning
- **Problems with lists:**
 - Using vague terms/traits:
 - **Why is it a problem?**
 - Ignoring non-psychological problems
 - CL has "solved" it (?)
 - **DO:** describe symptoms
 - **DO:** look for themes or relationships among problems

What should be on Mike's problem list?

- #1:
- #2:
- #3:
- #4:
- #5:

David Tolin (Doing CBT, 2016) Automatic + Semi-Automatic cognition

- **Automatic thoughts** > trigger mood congruent attention + recall
- **Intermediate beliefs** (Semi-Automatic)
 - Cognitive distortions, rules, interpretations
- **Core beliefs/schemas** (Semi-Automatic) > may trigger **compensatory strategies** (Young et al., 2006):
 - Maintenance/Surrender (do the usual...)
 - Avoidance/Escape (avoid your triggers)
 - Compensation/Counter-Attack (do the opposite)



Why behavior is so important

- **World responds to what we say/do, not what we think**
- **Guideline #1:** Do better in order to feel better
- **Guideline #2:** Do the healthy opposite:
 - Teaches new coping skills
 - Disconfirms one's (maladaptive) beliefs
- **Guideline #3:** Avoid avoidance:
 - Short-term gain may enable long(er) term pain

ESTs: Lack of response

- **40% of CLs are in Pre-Contemplation stage** (Prochaska et al., 2014)
- **Driessen et al., 2013:** 16 sessions of CBT vs. psychodynamic therapy:
 - No differences on any outcome measures
 - Average 22% remission
- **Friborg & Johnsen (2017):** results of CBT for unipolar depression declined over time

cbtscience/training/resources: Lack of progress worksheet

- **Sample "lack of progress" factors:**
 - Relationship is weak, problematic
 - Little is known about treating CL's disorder
 - Goals are unrealistic (or we disagree on them)
 - Treatment dose is not meeting CL's needs OR
 - CL needs adjunct (or different) treatment
 - My own/CL's behaviors are interfering w/treatment
 - Substance use is interfering with treatment
 - CL has high social strain or lacks social supports

David Burns:

Outcome + Process Resistance

- **Outcome:** CL resists due to “magical thinking,” i.e., superstitious beliefs about treatment outcomes:
 - My anxiety protects me from something worse
 - My depression is the price I must pay for my sins
- **Process:** CL resists interventions due to magical thinking:
 - Exposure treatment resistance: My anxiety protects me from X

Beck: Therapy interfering beliefs

- **If I try + solve problems >>>**
 - I'll fail OR have to become more responsible
 - It means my TH is controlling me + I'm weak
- **If I get better, my life will get worse. WHY?**
 - CL fears not meeting others' (new) expectations
 - Loss of social support or enabling relationships
 - Facing life challenges directly: may lose your disability, lose your therapist, etc.

Other treatment challenges (Ledley, et al., 2010)

- CL thinks s/he must discuss the past in order to get better
- CL thinks her/his problems are biologically determined
- CL thinks that CBT may not work for her/him
- Are CL's meds interfering with treatment?
 - CL's attributions for change when taking meds?

Integrating CBT + MI for anxiety (Randall & McNeil, 2016)

- **CBT elements consistent with MI:**
 - Problem-oriented > clear change targets
 - Highly collaborative relationship
 - Case formulation used to guide active treatment planning
 - Focus on skills + behavior change

Enhancing Motivation: Key MI processes

- **Engaging: Solid relational foundation**
 - Accurate empathy
 - **OARS** to understand ambivalence
 - Avoid the “righting” reflex (expert trap)
- **Focusing: Guide CL to a key change target:**
 - Identify behavior about which CL feels ambivalent
 - What's important to you? What could get in the way?

Enhancing Motivation: Key MI processes

- **Evoking: Draw out CL's reasons for change:**
 - Listen for “change talk” (vs. sustain talk)
 - Selectively reinforce + summarize change talk
 - Elicit-Provide-Elicit
- **Planning: Bridge to change:**
 - Selectively reinforce commitment language
 - Determine readiness for change + assist with specific change plans

Randall & McNeil (2016): Combining MI + CBT

- **Six case studies/uncontrolled trials + 11 RCTs:**
- *Treatment initiation and engagement can be improved by adding MI as an adjunct to CBT... (p. 308)*
- MI may improve:
 - Readiness to change
 - Treatment acceptance + commitment

Preparatory change talk: DARN

- **Desire to change**
- **Ability to change** (CL has self-efficacy)
- **Reasons to change** (the “pros” of changing)
- **Need to give up the status quo** (not changing)

Change Talk: CAT

- **Commitment:** stated intention to change:
 - I’m going to do X.....I plan to...
- **Activation:** leaning into change (I’ll try to...)
- **Taking steps: describe specific actions:**
 - I’ll walk every night after dinner...
 - I’ll try to (**activation**) eat 5 servings of fruit daily (**taking steps**)
 - I’ll limit myself to one drink (**taking steps**)

Elicit-Provide-Elicit

- **Elicit:** ask permission to make a suggestion
- **Provide:** suggest in **non-personal** ways:
 - **NOT** “You should do X”
 - **BUT:** “This has helped some CLs...”
- **Elicit:** explore CL’s reaction w/open ?s

Maximizing exposure treatment (Craske et al., 2014)

- **Fears may return after exposure treatment due to:**
 - Lack of repeated exposures after treatment ends OR
 - Phobic stimulus is encountered in a different context (EX: social phobia returns after a new social rejection)

Maximizing inhibitory learning (Craske et al., 2014)

- | | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> • Strategy • Expectancy violation • Deepened extinction • Reinforced extinction • Variability • Remove safety behaviors | <ul style="list-style-type: none"> • Catch phrase • Test it out • Combine it (2 cues) • Face your fear: present US in some trials • Vary it: vary stimuli + contexts • Throw it out: remove safety signals, acts |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

Strategies for maximizing inhibitory learning

- **Strategy**
 - Attentional focus
 - Affect labeling
 - Mental reinstatement or retrieval cues
- **Catch Phrase**
 - **Stay with it:** focus on CS during exposure
 - **Talk it out:** ask CL to describe emotions during exposure
 - **Bring it back:** use cue during extinction OR reinstate other successful exposures

CBT + Acceptance and Commitment Therapy

- Both address emotional self-regulation:
 - CBT uses **antecedent-focused** strategies
 - ACT uses **response-focused** strategies
- **Cognitive restructuring:** does it imply that your thoughts are “wrong”?
- **Heimberg + Ritter (2008):** CBT also promotes “cognitive defusion.”
 - Thoughts are “hypotheses” to test, NOT facts

Trans-theoretical approach: Stages of change

- **Prochaska et al (2014):** % CLs per stage >
 - Pre-Contemplation: 40-45%
 - Contemplation: 35-40%
 - Prepared for Action: about 20%
- Can double % taking action if progress one stage in a month

The right things (processes) at the right time (stages)

- **Rosen (2000):** 47 study meta-analysis:
 - Effect sizes of .70-.80 for use of different processes at different stages
 - Change processes for experiential, cognitive, psychoanalytic > best for Pre-Contemplation + Contemplation stages
 - Change processes for behavioral approaches: best for Action + Maintenance stages

Assessing stage of change (Prochaska et al., 2013)

- *Do you think behavior X is a problem for you now?*
 - If yes > Contemplation, Preparation, Action
 - If no > Maintenance, Pre-Contemplation
- *When do you intend to change behavior X?*
 - If not soon > Contemplation
 - If within next month > Preparation
 - If now > Action

References

- Arch JJ. & Craske, M.G. (2009) First line treatment: A critical appraisal of cognitive-behavioral therapy developments and alternatives. *Psychiatric Clinics of North America*, 32, 527-547. doi: 10.1061/j.psc.2009.05.001
- Balon, I.; Lejuez, C.W.; Hoffer, M. & Blanco, C. (2016). Integrating Motivational Interviewing and brief behavioral activation therapy: Theoretical and practical considerations. *Cognitive and Behavioral Practice*, 23, 205-220.
- Beck, J.B. (2005) *Cognitive therapy for challenging problems: What to do when the basics won't work*. New York: Guilford Press.
- Beck, J.B. (2011). *Cognitive behavior therapy: Basics and beyond*. (2nd edition) New York: Guilford Press.

References

Craske, M. G.; Treanor, M.; Conway, C.C.; Zbozinek, T. & Vervliet, B. (2014). Maximizing exposure therapy: An inhibitory learning approach. *Behavior Research and Therapy*, 58, 10-23.

Cristea, I.A; Stefan, S.; Karyotaki, K.; David, D.; Hollon, S. D. & Cuijpers, P. (2017). The effects of cognitive behavioral therapy are not systematically falling: A revision of Johnsen and Friborg (2015). *Psychological Bulletin* 143, 3, 326-340.

References

- Driessen, E. & Hollon, S.D. (2010). Cognitive behavioral therapy for mood disorders: Efficacy, moderators and mediators. *Psychiatric Clinics of North America*, 33, 537-555.

References

- Driessen, E.; Van, H.L.; Din, F. J.; Peen, J.; Kool, S.; Westra, D.; Henderickson, M.; Schoevers, R.A.; Cuijpers, P.; Twisk, J.W.; & Dekke, J.J. (2013). The efficacy of cognitive-behavior therapy and psychodynamic therapy in the outpatient treatment of major depression: A randomized clinical trial. *American Journal of Psychiatry*, 170, 1041-1050. doi: 10.1176/appi.ajp.2013.12070899

References

- Frank, E. & Levenson, J. (2010). *Interpersonal psychotherapy*. Washington, D.C.: American Psychological Association.
- Friborg, O. & Johnsen, T. J. (2017). The effect of cognitive-behavioral therapy as an antidepressive treatment is falling: Reply to Ljótsson et al. (2017) and Cristea et al. (2017). *Psychological Bulletin* 143, 3, 341-345.

References

- Heimberg, R.G. & Ritter, M.R. (2008). Cognitive-behavioral therapy versus Acceptance and Commitment therapy for the anxiety disorders: Two approaches with much to offer. *Clinical Psychology: Science and practice*, 15, 296-298.
- Knapp, S. & Gavazzi, J. (July/August 2017). What is new in psychotherapy processes and outcomes? *The Pennsylvania Psychologist*, 77, 5-6.

References

- Knapp, S. & Gavazzi, J. (July/August 2017). The changing evidence base for specific treatments. *The Pennsylvania Psychologist*, 77, 9-10.
- Leahy, R.L. (2001). *Overcoming resistance in cognitive therapy*. New York: Guilford Press.
- Leahy, R.L. (2003) *Cognitive therapy techniques: A practitioner's guide*. New York: Guilford.

References

- Ledley, D.R.; Marx, B.P.; and Heimberg, R.G. (2010). *Making cognitive-behavioral therapy work: Clinical process for new practitioners*. New York: Guilford Press.
- Miller, W. R. & Rollnick, S. R. (2012) *Motivational Interviewing: Helping people change* (3rd ed.). New York: Guilford press.

References

- Lowell, A. & Markowitz, J.C. (2017). You mean I have to talk about feelings? One CBT therapist's experience with Interpersonal Psychotherapy. *The Behavior Therapist*, 40, 181-185.
- Luty, S.E.; Carter, J.D.; McKenzie J.M.; Rae, A.M.; Frampton, C.M.; Mulder, R.D. & Joyce, P.R. (2007). Randomised controlled trial for cognitive behavioural therapy and interpersonal therapy for depression. *British Journal of Psychiatry*, 190, 496-502. doi: 10.1192/bjpp.bp.106.024729

References

- Newman, C. (1994). Understanding client resistance: Methods for enhancing motivation to change. *Cognitive Therapy and Research*, 1, 47-69.
- Nezu, A.M.; Nezu, C. M. & Lombardo, E. (2004). *Cognitive-behavioral case formulation and treatment design: A problem-solving approach*. New York: Springer Publishing.

References

- Norcross, J.C & Wampold, B.E. (2011). Evidence-based therapy relationships: Research conclusions and clinical practices. *Psychotherapy*, 48, 98-102.
- Persons et al, (2015). Developing + using a case formulation in cognitive-behavior therapy. *Journal of Psychology and Psychotherapy*, 5: 179. doi: 10.4172/2161-0487.1000179.

References

- Persons, J. ; Tompkins, M.A. & Davidson, J. (2000) *Cognitive-behavior therapy for depression: Individualized case formulation and treatment planning* (DVD). Washington, D.C.: American Psychological Association.
- Prochaska, J.O. and Norcross, J.C. (2014). *Systems of psychotherapy: A transtheoretical approach*. (8th ed.) New York: Cengage.

References

- Prochaska, J.O. and Norcross, J.C.; & DiClemente, C.C. (2013) Applying the stages of change. *Psychotherapy in Australia*, 19, 10-15.
- Randall, C.L. & McNeil, D.W. (2017). Motivational Interviewing as an adjunct to Cognitive Behavior Therapy for anxiety disorders: A critical review of the literature. *Cognitive and Behavioral Practice*, 24, 296-311.

References

- Rosen, C.S. (2000). Is the sequencing of change processes by stage consistent across health problems? *Health Psychology, 19*, 593-604.
- Simpson, H.B. & Zuckoff, A. (2011). Using Motivational Interviewing to enhance treatment outcome in people with obsessive-compulsive disorder. *Cognitive and Behavioral Practice, 18*, 28-37.

References

- Tolin, D. F. (2016) *Doing CBT: A comprehensive guide to working with behaviors, thoughts and emotions*. New York: Guilford Press.
- Young, J. E.; Klosko, J.S. & Weishaar, M.E. (2006) *Schema therapy: A practitioner's guide*. New York: Guilford Press.