

CIRCULAR OF INFORMATION

FOR THE USE OF CELLULAR THERAPY PRODUCTS

This circular was prepared jointly by the AABB, America's Blood Centers, the American Association of Tissue Banks, the American Red Cross, the American Society for Apheresis, the American Society for Blood and Marrow Transplantation, the College of American Pathologists, the Cord Blood Association, the Foundation for the Accreditation of Cellular Therapy, ICCBBA, the International Society for Cellular Therapy, the Joint Accreditation Committee of ISCT and EBMT, the National Marrow Donor Program, and Netcord. Federal law prohibits dispensing the cellular therapy products described in this circular without a prescription.

Contact Information

If you want to learn more about cellular therapy, contact any of the following co-sponsors of this publication.

AABB

www.aabcct.org

American Association of Tissue Banks (AATB)

www.aatb.org

American Red Cross (ARC)

www.redcross.org

**American Society for Blood and Marrow
Transplantation (ASBMT)**

www.asbmt.org

American Society for Apheresis (ASFA)

www.apheresis.org

America's Blood Centers (ABC)

www.americasblood.org

College of American Pathologists (CAP)

www.cap.org

Cord Blood Association (CBA)

www.cb-association.org/

**Foundation for the Accreditation of
Cellular Therapy (FACT)**

www.factwebsite.org

ICCBBA

www.iccbba.org

International NetCord Foundation

www.netcord.org

International Society for Cellular Therapy (ISCT)

www.celltherapysociety.org

JACIE Accreditation Office

www.jacie.org

National Marrow Donor Program (NMDP)

www.nmdp.org

Table of Contents

Contact Information	ii
Notice to All Users	1
General Information	1
Donors	2
Cellular Therapy Product Sources	3
HPC, Marrow	3
HPC, Apheresis	3
HPC, Cord Blood	3
Nucleated Cell Preparations	3
MNC, Apheresis	3
NC, Cord Blood	4
NC, Whole Blood	4
NC, Marrow	4
Cellular Therapy Product Descriptions	4
Actions	4
Indications	4
Contraindications	4
HPC, (Plasma reduced) Products	5
HPC, (RBC reduced) Products	5
HPC, (Buffy coat enriched) Products	5
HPC, (Mononuclear cell enriched) Products	5
HPC, Cryopreserved Products.....	5
HPC, (CD34 enriched) Products	6
Other Cellular Products	6
MNC, Apheresis; NC, Whole Blood; NC, Marrow	6
Instructions for Storage and Administration of Cellular Therapy Products	6
Dosage and Administration	7
Storage	7
Noncryopreserved Cellular Therapy Products	7
Cryopreserved Cellular Therapy Products	7
Cellular Therapy Product Labeling and Supporting Documents	8
Biohazard and Warning Labels	8
Side Effects and Hazards	9
Immunologic Complications, Immediate	9
Immunologic Complications, Delayed.....	10
Nonimmunologic Complications	11
Reporting of Adverse Reactions	14
References	19
Specific Product Information	20

Notice to All Users

The *Circular of Information for the Use of Cellular Therapy Products* (hereafter referred to as the *Circular*) is an extension of container labels, as the space on those labels is limited.¹ The focus of this *Circular* is restricted to unlicensed cellular therapy products that are minimally manipulated. These unlicensed products can be: hematopoietic progenitor cells (HPCs), leukocytes and other cells derived from bone marrow, umbilical cord blood, or cellular products collected by apheresis. This *Circular* does not apply to products that have already received a license as a cellular therapy product in the United States. Per requirements of other national competent authorities, cellular therapy products may be designated as licensed biological products, medical devices, or advanced therapy medicinal products. Principles expressed here may also be applied to other cellular therapy products. Cellular therapy products are biological products that contain living human cells and are intended for use in patient treatment. Professional judgment based on clinical evaluation determines the selection of products, dosage, rate of administration, and decisions in situations not covered in this general statement.

WARNING: Because cellular therapy products are derived from human blood or tissues, they may carry a risk of transmitting infectious agents, including bacteria, viruses, fungi, protozoa, and prions. Donor screening and testing procedures are in place to minimize the risk of transmitting such infections but cannot eliminate this risk. Transmission of malignant disease has been reported. Also, serious life-threatening septic and toxic reactions can result from administration of products containing bacterial toxins. In addition, cellular therapy products may contain certain immunizing substances other than those indicated on the label, such as red cells, mature white cells, platelets, and plasma proteins. Therefore, this *Circular*, in whole or in part, cannot be considered or interpreted as an expressed or implied warranty of the safety or fitness of the described products even when they are used for their intended purpose. Attention to the specific indications for cellular therapy products is needed to prevent inappropriate administration.

This *Circular* addresses some of the applicable regulations established by regulatory/competent authorities such as the Food and Drug Administration (FDA), the Health Resources and Services Administration (HRSA), and Directive 2004/23/EC (and other European Commission directives) of the European Parliament and the Council of the European Union (EU).²⁻⁷ This *Circular* is not a comprehensive reference for applicable regulations.

The nomenclature used throughout this *Circular* is consistent with *ISBT 128* terminology and was current at the time of publication.⁸⁻¹⁰ However, acronyms such as HPC(CB), MNC(A), and HPC(M) are used only as abbreviations and are not intended to be used on the full product labels. (Check ICCBBA's standard terminology⁸ for appropriate usage of acronyms.) Users of this *Circular* should confirm that the terminology is still in effect before labeling and distributing a cellular therapy product for patient use.

General Information

This *Circular* was prepared by the Circular of Information for Cellular Therapy Products Task Force, consisting of representatives from the AABB, the American Association of Tissue Banks (AATB), the American Red Cross (ARC), the American Society for Blood and Marrow Transplantation (ASBMT), the American Society for Apheresis (ASFA), America's Blood Centers (ABC), the College of American Pathologists (CAP), the Foundation for the Accreditation of Cellular Therapy (FACT), ICCBBA, the International Society for Cellular Therapy (ISCT), the National Marrow Donor Program (NMDP), the Joint Accreditation Committee of ISCT and EBMT (JACIE), and NetCord. The text of this document has been approved by the board of directors of each of these organizations. Representatives from the FDA and HRSA participated in the deliberations of this task force.

This *Circular* is intended to provide general information to those who administer cellular therapy products, and serves as an extension and enhancement of the label found on the cellular therapy product. The task force has chosen to describe only those cellular therapy products that are most frequently used in clinical practice. **Not all cellular therapy products are described in this *Circular*.**

In order to address other cellular therapy products that are not listed in the *Circular*, this document is designed with a section of blank pages at the end to allow for inclusion of facility-specific information. It is important for users of this document to examine this section of the *Circular* for any additional information provided by the distributing facility and/or the manufacturer of the cellular therapy product. The portion preceding this section of the document cannot be changed.

This *Circular* is intended to be used by facilities based in different countries. The task force has made a concerted effort to accommodate both US and EU requirements in the document text. However, the regulatory approaches to cellular therapy products in the United States and the EU, as well as in other countries, differ in some aspects. Users should consult the appropriate regulatory authority for specific requirements related to their facility.

For investigational products manufactured and administered in the United States, an FDA-approved investigational new drug (IND) application or an investigational device exemption (IDE) is required. For investigational products manufactured and administered outside the United States, other local regulations apply. The relevant clinical protocol should be consulted for information regarding the indications for use, specific details for the administration of the product, and any expected toxicities. For corporate-sponsored or multicenter clinical trials, the indications and administration and toxicity information can also be found in the investigator's brochure.

Donors

Cellular therapy products described in this *Circular* have been collected from human donors for autologous or allogeneic administration. Autologous HPC collection usually occurs after mobilization of the donor's stem and progenitor cells with growth factors, chemotherapy, or both. Donors of other cellular therapy products may or may not require stimulation by growth factors, depending on the protocol employed. Allogeneic HPC collection usually occurs after mobilization with growth factors alone. Certain products such as HPCs from marrow [HPC, Marrow; or HPC(M)] and mononuclear cells from apheresis [MNC, Apheresis; or MNC(A)] are usually collected from donors who are not mobilized.

US federal regulations do not require testing of autologous donors for transmissible agents. However, the voluntary accrediting organizations (eg, AABB, FACT-JACIE) and some states or regions may require additional testing and/or testing of autologous donors. Allogeneic donors are screened through the use of questions designed to detect risk factors for infectious diseases transmissible by the cellular therapy product. Allogeneic donors are also tested for transmissible infectious diseases. (See Tables 1A and 1B.) The questions are based on donor screening requirements promulgated by regulatory agencies and criteria set forth by standard-setting organizations. A donor questionnaire and accompanying donor screening materials* have been developed for cellular therapy products and cord blood donors. The provision of truthful and accurate information by a donor during health/risk assessment is essential for the exclusion of donors whose cellular therapy products may transmit diseases to recipients.

Some allogeneic donors may not meet all the requirements; however, because of the patient's clinical circumstances, they may be approved for donation. In such situations, information regarding requirements that the donor has not met is included in the summary of records/information provided to the transplant center. The cellular therapy products from such donors are also labeled accordingly. (See Table 2.) Cellular therapy products from a donor with abnormal screening and/or test results may be administered to a recipient if the recipient has been advised of the risk, the recipient's physician has authorized the use of the product, and the product is appropriately labeled.

*An example of such a questionnaire, called the uniform donor questionnaire, has been prepared with input from the AABB, AATB, ASFA, and FACT, and can be accessed on the AABB website (www.aabb.org) under "Programs & Services" > "Transfusion Medicine" > "Donor History Questionnaires."

Cellular Therapy Product Sources

HPC, Marrow

HPC, Marrow [HPC(M)] preparations are collected as a source of HPCs. They are obtained by multiple needle aspirations from the posterior iliac crests of an autologous or allogeneic donor. The marrow is placed in a sterile container with an electrolyte solution and an appropriate anticoagulant. The cell suspension is passed through sterile filters to remove fat, bone particles, and cellular debris. The volume of HPC(M) products varies and may range from 100 mL to 2000 mL. Marrow contains mature red cells, white cells, platelets, committed progenitors of all lineages, mast cells, fat cells, plasma cells, and pluripotent hematopoietic cells. Some of these cells are capable of reconstituting the hematologic and lymphoid systems of an autologous or allogeneic recipient. These cells are usually processed before infusion but are sometimes infused in an unmodified state. The most common modifications of allogeneic HPC(M) products are reduction of the volume of ABO-incompatible red cells, removal of ABO-incompatible plasma, selection of CD34+ progenitor cells, and removal of donor T lymphocytes. The most common modification of autologous HPC(M) products is to reduce the volume by removing plasma and red cells before cryopreservation.

HPC, Apheresis

HPC, Apheresis [HPC(A)] preparations are collected as a source of HPCs obtained from the peripheral blood using an apheresis technology, usually after recombinant hematopoietic growth factor administration or other agents. Autologous donors may also have undergone chemotherapy mobilization. Allogeneic HPC(A) collections are frequently infused without further processing. Additional processing of allogeneic HPC(A) products includes reduction or removal of ABO-incompatible red cells, removal of ABO-incompatible plasma, selection of CD34+ progenitor cells, or removal of donor T lymphocytes. Common types of additional processing of autologous HPC(A) products are reduction of volume by removing plasma before cryopreservation and selection of CD34+ progenitor cells. HPC(A) products may be thawed and washed to remove dimethyl sulfoxide (DMSO).

HPC, Cord Blood

HPC, Cord Blood [HPC(CB)] preparations are collected as a source of HPCs obtained from the umbilical cord during the third stage of labor or after delivery of the placenta. After thorough cleansing of the cord, the blood is collected by gravity drainage into standard collection bags containing anticoagulant. Before cryopreservation, cord blood collections are usually processed by red cell and plasma reduction. HPC(CB) products are typically stored with final 10% DMSO cryoprotectant in bags with integral segments designed to be a source of material for identity and potency testing. Frozen cord blood products are transported to the transplant center before patient conditioning begins and are typically thawed using a wash or reconstitution method before infusion. HPC(CB) products that are not red cell reduced should be washed or diluted to lessen the potential effects of hemolysate.

Nucleated Cell Preparations

MNC, Apheresis

MNC, Apheresis [MNC(A)] preparations contain nucleated cells collected from the peripheral blood by an apheresis procedure and are intended for clinical use other than as HPCs. Autologous MNC(A) collections are generally further processed. Allogeneic MNC(A) collections are most commonly used as donor lymphocyte infusions (DLIs). The dose for MNC(A) is determined by institutional policies and is usually based on the number of T cells (eg, CD3+ cells), nucleated cells, or mononuclear cells.

NC, Cord Blood

NC, Cord Blood [NC(CB)] preparations are collected as a source of nucleated cells obtained from the umbilical cord during the third stage of labor or after delivery of the placenta and are intended for clinical use other than as HPCs.

NC, Whole Blood

NC, Whole Blood [NC(WB)] preparations contain nucleated cells collected as peripheral whole blood and are intended for clinical use other than as HPCs.

NC, Marrow

NC, Marrow [NC(M)] preparations contain nucleated cells collected from bone marrow and are intended for clinical use other than as HPCs.

Cellular Therapy Product Descriptions

Cellular therapy products consist of somatic-cell-based products [eg, HPC(A), HPC(M), HPC(CB), MNC(A), NC(WB)] that are collected or procured from the donor and intended for manipulation and/or administration to the patient.

HPC products contain hematopoietic stem and progenitor cells capable of providing hematopoietic and immune reconstitution after myeloablative or nonmyeloablative preparative regimens. The products contain pluripotent and lineage-committed hematopoietic progenitors as well as lymphocytes.

Actions

HPCs administered intravenously migrate to the marrow, where they divide and mature. The mature cells are released into the bloodstream, restoring blood counts and immunity.

The time from administration of HPCs to recovery of adequate or normal blood counts is variable. Allogeneic transplantation sometimes induces a graft-vs-tumor effect that is beneficial in recipients who receive a transplant for treatment of malignancies.

Allogeneic cellular therapy products may also be used to provide additional donor lymphocytes to enhance a graft-vs-leukemia effect. Other applications of cellular therapy products may have different potential mechanisms of action depending on the cell type and clinical setting.

Indications

Allogeneic HPC products are intended to provide hematopoietic reconstitution after myeloablative or nonmyeloablative preparative regimens for a wide range of disease states. For patients with certain malignancies, the product is also intended to provide immune reconstitution and immune-mediated therapy. Autologous HPCs are collected and used following myeloablative or myelotoxic therapy to enhance hematopoietic reconstitution. The therapy is intended to treat the patient's underlying malignancy, and autologous HPC products are administered to minimize morbidity and mortality caused by the myelotoxic effects of the therapy. Additional applications may be used as indicated in clinical trials and research protocols.

Contraindications

MNC(A) and NC(WB) are generally contraindicated for patients experiencing severe graft-vs-host disease (GVHD). Institutional policies and protocols and federal regulations dictate specific contraindications for cellular therapy product administration. Additional information regarding contraindications may be included at the end of this document, if provided by the distributing facility.

The following section provides common cellular therapy product descriptions in the product description format consistent with *ISBT 128* information and labeling standards.

HPCs contain self-renewing or multipotential stem cells capable of maturing into any hematopoietic lineage, lineage-restricted pluripotent progenitor cells, and committed progenitor cells. They may be collected from bone marrow [HPC(M)], peripheral blood with or without prior mobilization [HPC(A)], whole blood with mobilization (HPC, Whole Blood), or placental/umbilical cord blood [HPC(CB)]. They may then be subjected to volume reduction or further manipulations. (See below.)

HPC, (PLASMA REDUCED) PRODUCTS

HPC, APHERESIS	HPC, CORD BLOOD
Plasma Reduced	Plasma Reduced

HPC, MARROW
Plasma Reduced

These products contain the cellular elements of the starting HPC collection(s) that remain after the bulk of the plasma is removed by centrifugation.

HPC, (RBC REDUCED) PRODUCTS

HPC, CORD BLOOD
Red Cells Reduced

HPC, MARROW
Red Cells Reduced

These are the HPCs remaining after the mature red cells have been reduced by sedimentation, centrifugation, or lysis.

HPC, (BUFFY COAT ENRICHED) PRODUCTS

HPC, CORD BLOOD
Buffy Coat Enriched

HPC, MARROW
Buffy Coat Enriched

The buffy coat is the portion of an HPC product containing the nucleated cells after the bulk of the plasma and mature red cells has been removed by sedimentation or centrifugation techniques.

HPC, (MONONUCLEAR CELL ENRICHED) PRODUCTS

HPC, CORD BLOOD
Mononuclear Cell Enriched

HPC, MARROW
Mononuclear Cell Enriched

These are primarily mononuclear cells that remain after the depletion of mature red cells, polymorphonuclear leukocytes, and plasma by separation of the cells on the basis of their density. This is achieved using devices or density gradient solutions.

HPC, CRYOPRESERVED PRODUCTS

HPC, APHERESIS	HPC, CORD BLOOD
CRYOPRESERVED	CRYOPRESERVED

HPC, MARROW
CRYOPRESERVED

These are HPCs that have been frozen using cryoprotectant solutions and containers suitable for the purpose.

HPC, (CD34 ENRICHED) PRODUCTS

HPC, APHERESIS
CD34 Enriched

HPC, CORD BLOOD
CD34 Enriched

HPC, MARROW
CD34 Enriched

These products contain the cellular elements of HPCs that have been enriched by selection of CD34+ cells.

Other Cellular Products

These are nucleated cells from any source (marrow, peripheral blood, whole blood, or umbilical cord/placental blood) and intended for clinical use other than as HPCs. They may be further categorized according to the specific subpopulations.

MNC, APHERESIS; NC, WHOLE BLOOD; NC, MARROW

These products are most frequently used for DLIs. They are usually collected from the HPC donor and contain a mixture of mature nucleated cells (eg, T and B lymphocytes, granulocytes), red cells, and plasma.

Instructions for Storage and Administration of Cellular Therapy Products

The following instructions pertain to cellular therapy products described in this *Circular*:

- All products must be maintained in a controlled environment and stored under appropriate conditions as described in FDA regulations and applicable AABB, AATB, FACT-JACIE, NetCord-FACT, or NMDP standards.¹²⁻¹⁶
 - Note:** If the administration of a cellular therapy product is delayed, the distributing/issuing facility should be contacted for instructions on proper storage of the product during the delay.
- Before administration of the product, it is critical to coordinate patient and product preparation to support timely product infusion according to the facility standard operating procedure. Infusion coordination may include confirmation of the number of containers and type of product (fresh or cryopreserved), verification of product infusion order, verification of consent for infusion, and verification of patency of intravenous access for infusion of the product.
- The intended recipient and the product container must be properly identified according to facility standard operating procedure before the product is administered.
- The product must be inspected for changes in the integrity of the container and product condition before administration. Any questions about the product should be directed to the facility distributing or issuing the product.
- Aseptic technique must be employed when handling and administering the product.
- Products must *not* be administered through a filter designed to remove leukocytes.
- Products may be filtered through a 170- to 260-micron filter designed to remove clots.
- Products should be mixed thoroughly before use.
- Products must *not* be irradiated.
- No medications or solutions may be added to or infused through the same tubing as products, with the exception of 0.9% Sodium Chloride, Injection (USP) or facility-approved solutions, as directed by the distributing facility. Periodic observation of the patient is required during and after administration to detect adverse reactions. Vital signs must be recorded at a minimum before and after administration, or more often, if required, by facility standard operating procedure.

- Sequence and timing of multiple product infusions should be performed according to the administering facility's standard operating procedures. Adequate time between product infusions should be allowed to permit assessment for adverse reactions.

Dosage and Administration

The minimum number of HPCs necessary for engraftment in a myeloablated recipient has not been established for all HPC sources. However, eligibility criteria for some protocols may dictate a minimum number of cells to be collected and/or infused. Some examples of cell types measured to determine HPC dosage are CD34+ cells, nucleated or mononuclear cells, and colony-forming units–granulocyte-macrophage (CFU-GM). The dose for MNC(A) or NC(WB) is determined by institutional policies and is usually based on the number of T cells, nucleated cells, or mononuclear cells. For specific dosage and administration of other cellular therapy products, the investigator's brochure or special instructions should be followed. Such information may be found at the end of this document, if provided by the distributing facility. Administration of any cellular therapy product should begin only after identification of the product(s) and the intended recipient according to institutional policies. Manufacturers may recommend that products be filtered using a 170- to 260-micron filter to remove clumps or aggregates. Some institutions may have specific policies regarding the use of these filters for cellular therapy products. (See facility-specific section at the end of this document.) Cellular therapy product infusion should begin slowly and with sufficient observation to detect symptoms and/or signs suggestive of acute immunologic or infectious complications. Thereafter, the rate of infusion may be as rapid as tolerated. The administration time will be determined by the total volume to be infused and by whether the cells are fresh or previously cryopreserved. If the thawed products have not been washed to remove DMSO, care should be taken not to exceed 1 mL of DMSO per kilogram of recipient weight per day administration (eg, 100 mL of a 10% solution contains 10 mL of DMSO).

Storage

Cellular therapy products may be transported for administration in a fresh or cryopreserved state. They may require either long-term or short-term storage before administration. Institutional policies and protocols dictate specific storage requirements for cellular therapy products. The recommended storage duration and temperature may be included in the product labeling and should indicate the cell dose, storage temperature, and duration of storage to ensure acceptable cell viability and function. If an expiration date has been defined, it should be included on the product label. Before infusion, products received for the treatment of a patient should be stored according to the instructions on the label or those supplied in accompanying documentation. If there is an unexpected delay in administration and the product must be held for infusion after the expiration period indicated on the label, if applicable, the distributing/issuing and/or local cell processing facility should be contacted for further handling and storage instructions.

Noncryopreserved Cellular Therapy Products

Fresh products may be transported from distant collection facilities or undergo short-term local storage before administration.

Cryopreserved Cellular Therapy Products

Cryopreserved products may be received and stored long-term according to the manufacturer's directions or by a validated method. These products may be thawed at the local cell processing laboratory, with or without additional processing, or thawed at the bedside immediately before administration. These products should be infused as soon as possible after thawing occurs.

Cellular Therapy Product Labeling and Supporting Documents

At the time of issue, cellular therapy products will have the following information either on the affixed product label, on an attached label, or in accompanying documentation:

- Proper name of the product, including an indication of any qualification or modification.
- Unique identifier.
- Approximate volume.
- Name and volume of anticoagulant or other additives.
- Date and end time of collection.
- Expiration date and time (if applicable).
- Recommended storage temperature.
- Identity and address of collection facility or donor registry.
- Identity and address of processing/distributing facility.
- Statements regarding transmission of infectious diseases.
- Statement indicating “Do Not Irradiate.”
- Biohazard or other warning label(s) (if applicable),
- Statements regarding recipient identification,
- Donor identifier and (if applicable) name.
- Recipient name and identifier.
- ABO group and Rh (D) type of donor or the ABO group and Rh (D) type of a cord blood product.
- Red cell compatibility testing results (if applicable).

Many products will be accompanied by additional records that are included to meet regulatory requirements. These accompanying records will include:

- A statement indicating whether the donor has been determined to be eligible or ineligible, or that the donor eligibility determination is incomplete.
- A summary of the records used to make the donor eligibility determination.
- Infectious disease testing results and supporting documents.
- For ineligible donors, a statement noting the reasons for ineligibility [21 CFR 1271.55(b)(4)].
- For products that are made available before the donor eligibility has been completed:
 - The results of any donor screening and testing that has been completed [21 CFR 1271.60(d)(2)(i) and (ii)].
 - A list of any screening and testing that has not yet been completed [21 CFR 1271.60(d)(2)(iii)].

International standards for nomenclature and labeling of cellular therapy products using *ISBT 128* have been determined by the International Cellular Therapy Coding and Labeling Advisory Group.⁸⁻¹⁰

Biohazard and Warning Labels

The application of biohazard and warning labels on the cellular therapy products summarized in Table 2 is defined by facility-specific policies and procedures. The FDA defines requirements for the use of biohazard and specific warning labels for products subject to the regulations as defined in 21 CFR 1271, implemented on May 25, 2005. As such, cellular therapy products subject to these FDA regulations require the use of these labels as specified by FDA for an “incomplete” or “ineligible” donor eligibility determination. Refer to 21 CFR 1271 for specific labeling guidance. Application of these labels extends outside the FDA-defined requirements, such as to HPC(M), based on voluntary adherence to professional industry standards and facility-specific guidance or other applicable laws. Questions about the interpretation of any label on a specific product should be directed to the facility distributing the product.

Side Effects and Hazards

Some side effects and complications may require reporting to a relevant national competent authority. See “Reporting of Adverse Reactions” later in this *Circular* for additional details.

The following side effects and hazards pertain to administration of cellular therapy products.

Immunologic Complications, Immediate

1. **Acute Hemolytic Reactions** can be a complication of cellular therapy product administration and can be caused by donor-recipient major or minor ABO or other blood group incompatibility. Acute hemolytic reactions may be immediate and occur up to 24 hours following infusion.

Signs and symptoms of acute hemolytic reactions may include one or more of the following:

- Chills.
- Fever.
- Headache.
- Burning sensation along the vein.
- Abnormal bleeding.
- Low back pain.
- Facial flushing.
- Chest pain; rapid, labored respirations.
- Tachycardia.
- Shock.
- Hemoglobinuria:
 - Development of a positive direct antiglobulin test (DAT).
 - Elevation of lactate dehydrogenase (LDH) or bilirubin.

Treatment:

- Measures to maintain or correct arterial blood pressure; correct coagulopathy, if present; and promote and maintain urine flow.

Prevention:

- Red cell reduction.
- Plasma reduction.

2. **Febrile Nonhemolytic Reactions** may reflect the action of cytokines, either present in the product or generated by recipient antibodies against infused white cells. These reactions occur more frequently in patients previously alloimmunized by transfusion or pregnancy.

Signs and symptoms of febrile nonhemolytic reactions include:

- Temperature elevation of 1 C (2 F) or more (shortly after or up to 2 hours following product administration and in the absence of another pyretic stimulus).
- Chills.

Treatment:

- Antipyretics.

Prevention:

- Antipyretics.

3. **Allergic/Anaphylactoid/Anaphylactic Reactions** are thought to be related to the presence of atopic substances capable of interacting with antibodies present in the donor or recipient plasma. In rare cases, anaphylaxis may occur. These reactions have typically been reported in IgA-deficient patients who have IgA-specific antibodies of the IgG and/or IgE class and who receive even small amounts of IgA-containing plasma. Allergic reactions to hydroxyethyl starch (HES) or DMSO used in cellular therapy product processing or cryopreservation may occur in sensitized patients.

Signs and symptoms of allergic reactions include:

- Urticaria (hives).
- Pruritus (itching).

- Bronchospasm and/or laryngospasm.
- Hypotension.
- Severe dyspnea.
- Facial, glottal, and/or laryngeal edema.
- Other symptoms such as facial burning and flushing, abdominal pain, nausea, vomiting, diaphoresis, diarrhea, and dizziness.

Treatment:

- Antihistamines.
- In severe cases, fluids, epinephrine, and/or steroids.

Prevention:

- Premedication with antihistamines is sometimes used to mitigate mild reactions.
- Washing of products can help prevent symptoms, but this procedure is usually reserved for patients with a history of severe/anaphylactic reactions.

4. Transfusion-Related Acute Lung Injury (TRALI) occurs when an acutely increased permeability of the pulmonary microcirculation allows massive leakage of fluids and protein into the alveolar spaces and interstitium. In many cases, the occurrence of TRALI is associated with the presence of leukocyte antibodies (eg, anti-HLA) in the donor or recipient. As such, these reactions are rare in recipients of HLA-matched products.

In the absence of evidence for another cause of pulmonary compromise, signs and symptoms of TRALI include:

- Acute respiratory distress within 6 hours of administration.
- Hypoxemia.
- Bilateral pulmonary infiltrates on frontal chest x-ray.

Treatment:

- Respiratory support.

Prevention:

- Plasma-reduction or washing can help reduce the risk of TRALI in the setting of a graft with known anti-HLA or antibody to human neutrophil antigen (HNA), but these procedures are rarely performed for this indication.

Immunologic Complications, Delayed

1. **Alloimmunization to Antigens** of red cells, white cells, platelets, or plasma proteins may occur after infusion of cellular products. Primary immunization does not become apparent until days or weeks after the immunizing event and does not usually cause symptoms or physiologic changes. However, in patients who have developed alloantibodies, if blood or cellular therapy products that express the relevant antigens are subsequently administered, there may be accelerated removal of cellular elements from the circulation and/or systemic symptoms that may contribute to graft failure, red cell aplasia, and transfusion refractoriness.

Treatment:

- Selective use of blood components for transfusion support that are negative for the antigen recognized by the alloantibody.

2. **Delayed Hemolytic Reactions** may occur in two different allogeneic settings. Clinically significant antibodies to red cell antigens in previously alloimmunized patients are usually detected by pretransfusion testing. Occasionally, however, levels may diminish to below the limits of detection. In these cases, antigens on transfused red cells can stimulate anamnestic production of antibody. The antibody levels may reach a significant circulating level while the transfused red cells are still present in the circulation, leading to hemolysis. The usual time frame for reappearance of antibody is 2 to 14 days after product administration. Delayed hemolysis may also occur in recipients who receive plasma-incompatible transplants, whether in regard to ABO antigens or to other red cell antigens. In

this setting, the infused donor's B lymphocytes may produce antibodies to red cell antigens, thus destroying the recipient's own remaining red cells in the 1 to 3 weeks after HPC product administration. This reaction may be sudden, severe, and life threatening, so at-risk recipients should be monitored for this occurrence.

Signs and symptoms of delayed hemolytic reactions may include:

- Unexplained fever.
- Unexplained decrease in hemoglobin/hematocrit.
- Mild jaundice.
- Development of a positive DAT.
- Elevation of LDH or bilirubin.
- Hemoglobinemia and hemoglobinuria (rare).
- Symptoms of acute intravascular hemolysis (rare).

Treatment:

- Use of antigen-negative red cells if transfusion is needed.
- More severe cases may require treatment similar to an acute hemolytic reaction and more rapid antigen-negative red cell replacement.

Prevention:

- Providing red cells after transplantation that are ABO compatible with the donor and recipient.

3. ***Graft-vs-Host Disease*** (GVHD) is an extremely serious condition that occurs frequently in recipients of allogeneic cellular therapy products.¹⁷ GVHD occurs when viable T lymphocytes in the infused product engraft and react against tissue antigens in the recipient.

Signs and symptoms:

- Wide variety of immune-mediated tissue and organ damage.

Treatment:

- Posttransplant immunosuppression, according to institutional guidelines and policies.

Prevention:

- Use of optimally matched HLA-compatible donor grafts.
- Incorporation of immune suppression into the transplant regimen.

Nonimmunologic Complications

1. ***DMSO Toxicity*** is the most common complication of cryopreserved product administration. DMSO is a cryoprotectant used to cryopreserve cellular therapy products. Side effects and symptoms are generally associated with histamine release.

Signs and symptoms:

- Coughing.
- Flushing.
- Rash.
- Chest tightness and wheezing.
- Nausea and vomiting.
- Cardiovascular instability.

Treatment:

- Slowing the rate of infusion.
- Medicating with antihistamines.
- Treating symptoms.

Prevention:

- Prophylactic antihistamine therapy.
- Decreased rate of administration.
- Providing hard candy to prevent nausea caused by the odor and/or taste.

- Removing DMSO from the product by washing the cells before administration; while this may reduce the risk of symptoms, it is not generally required given typically mild reactions, and it may result in unintended cell loss.

2. **Septic Infusion Reactions** may result from bacterial contamination of cellular therapy products, but they rarely cause acute, severe, or life-threatening effects. Prompt recognition of a possible septic reaction is essential. The onset of high fever (>2 C rise in temperature) during or immediately after product administration should suggest the possibility of bacterial contamination and/or the presence of endotoxin in the product.

Possible signs and symptoms of septic infusion reactions include:

- Fever with chills.
- Severe hypotension.
- Dry, flushed skin.
- Pain in abdomen and extremities.
- Vomiting.
- Bloody diarrhea.

Treatment:

- Prompt and appropriate use of antimicrobial agents with modification based on evaluation of blood culture results from the patient and the product when available.

Prevention:

- Appropriate aseptic technique during all aspects of product collection, manufacturing, and infusion.
- Appropriate antibiotic prophylaxis should be considered when using nonconforming products with positive culture results according to institutional protocol and relevant national competent authorities.

3. **Fat Emboli**, small fat droplets in marrow products, may block capillary perfusion and cause respiratory distress.

Signs and symptoms of fat emboli syndrome (FES) include:

- Dyspnea.
- Hypoxia.
- Tightness of the chest.
- Coughing.
- Petechiae.
- Confusion (mental status change).

Treatment:

- Supplemental oxygen therapy.
- Ventilation as needed.
- Corticosteroids, including methylprednisolone, which have reduced posttraumatic hypoxemia believed to be due to FES.

Prevention:

- Routine filtering of bone marrow products with 170- to 260-micron filters before infusion.

4. **Transmission of Infectious Disease and/or Disease Agents** may occur because cellular therapy products are collected from human blood and/or tissues. Disease may be caused by known or unknown agents. Donor selection criteria, screening, and testing are designed to minimize the potential risk of disease transmission. These procedures aim to identify potential donors with increased risk of infection with human immunodeficiency virus (HIV), human T-cell lymphotropic virus (HTLV), hepatitis B virus (HBV), hepatitis C virus (HCV), and syphilis, as well as other agents. (See “Donors” section.) These measures do not totally eliminate the risk of transmitting these agents. Cytomegalovirus (CMV) may, unpredictably, be present in white-cell-containing products from donors previously infected with this virus, which can persist lifelong despite the presence of serum antibodies. Up to 70% of donors

may be CMV-seropositive. Transmission of CMV may be of concern in immunocompromised transplant recipients if they are CMV-seronegative. Administering CMV-seronegative cellular therapy products reduces the risk of CMV transmission. For some infectious agents, there are no routine tests to predict or prevent disease transmission.

Treatment:

- Based on implicated infectious agent.

Prevention:

- Minimize by robust screening procedures, identification of infectious donors, and proper labeling.

5. ***Bleeding Due to Excessive Anticoagulation*** can occur if heparin or other anticoagulants were added to the product during collection and/or processing and remain in the cellular therapy product when administered.

Treatment:

- Anticoagulant specific.
- A reversal agent can be considered.

Prevention:

- Product/anticoagulant specific.
- Infusion rates may be adjusted depending on the clinical conditions; products may be washed when cell loss is not a concern.

6. ***Circulatory Overload*** leading to pulmonary edema can occur after infusion of excessive volumes or at excessively rapid rates. Pulmonary edema should be promptly and aggressively treated. In at-risk patients, the infusion of colloid preparations (including plasma products and the suspending plasma in cellular therapy products) should be reduced to a minimum.

Signs and symptoms of circulatory overload may include:

- Dyspnea.
- Peripheral edema.
- Rapid increase of blood pressure.

Treatment:

- Diuresis.

Prevention:

- Minimize the volume of colloidal preparations and, if appropriate, split or volume-reduce the product for infusion.
- Reduce the rate of administration.

7. ***Hypothermia*** is related to the temperature of the cellular therapy product and the rate of infusion and can be caused by rapid infusion of large volumes of cold products. Hypothermia carries a risk of cardiac arrhythmia or cardiac arrest. A blood warming device should not be used unless approved by the manufacturer of the cellular therapy product.

Treatment:

- Warm the patient.

Prevention:

- Decrease infusion rate when clinically appropriate.

8. ***Nonimmunologic Hemolysis*** can result from lysis of red cells in the product, which may occur at any time during processing, cryopreservation, thawing, and administration. This lysis may be caused by osmotic stress, mechanical injury, shear stress, coadministration with incompatible fluids, or intrinsic red cell abnormalities such as hemoglobinopathies or enzyme deficiencies. Some hemoglobinuria can be seen even with products containing only small amounts of free hemoglobin and does not necessarily indicate a reaction.

Signs and symptoms:

- May be the same as hemolytic reactions, either delayed or immediate.

Treatment:

- Same as treatment of hemolytic reactions.

Prevention:

- High levels of free hemoglobin can be removed by washing the product when clinically appropriate and using isotonic solutions during product preparation.
- Prevention relates to proper product handling during all steps of product collection, manufacturing, and administration.

Reporting of Adverse Reactions

Any adverse reaction that is defined as a suspected or proven unfavorable response to administration of cellular therapy products and is manifested by signs and symptoms (including microbial contamination of a product or suspected disease transmission during or after product administration) must be documented and reported in accordance with the facility's policies and/or applicable laws and regulations. At a minimum, any such event must be reported to the patient's physician and to the medical director of the facility that issued the product.

The reporting requirements vary based on the regulatory oversight required by the type of product and manufacturing process. The user must contact the manufacturing/distributing facility for specific requirements.

Entities involved in the manufacture of the product must be contacted in the investigation/reporting of an adverse reaction, as applicable.

Table 1A. US Minimal Requirements for Testing for Transmissible Agents in Cellular Therapy Products*[†]

Testing for Infectious Agents	Donors of HPC(M) and HPC(A)	Donors of HPC(CB) and NC(CB)	Donors of NC(WB), MNC(A), and NC(M)
Timing of specimen collection	Up to 30 days before or 7 days after collection	Up to 7 days before or after collection	Up to 7 days before or after collection
Human immunodeficiency virus, types 1 and 2 (HIV-1, HIV-2) [‡]	X	X (MS)	X
Hepatitis B virus (HBV)	X	X (MS)	X
Hepatitis C virus (HCV)	X	X (MS)	X
Human T-cell lymphotropic virus, types I and II (HTLV-I, HTLV-II)	X	X (MS)	X
Cytomegalovirus (CMV) (if allogeneic)	X	X (MS)	X
<i>Treponema pallidum</i> (syphilis)	X	X (MS)	X

*Testing is performed according to manufacturers' instructions using donor screening tests that have been licensed, approved, or cleared by the US Food and Drug Administration (FDA) for transmissible agents as defined by the FDA.¹¹ More than one test may need to be conducted to adequately and appropriately test for a single communicable disease agent or disease. Refer to the Center for Biologics Evaluation and Research website for donor eligibility guidance and a list of approved tests. FDA-licensed, -approved, or -cleared donor screening tests are available for West Nile Virus (WNV) and HBV nucleic acid testing (NAT) and *Trypanosoma cruzi*; testing may be implemented per facility-specific guidance prior to an FDA testing requirement. Additional tests for infectious transmissible agents may be required or performed.

[†]US federal regulations do not require testing of autologous donors for transmissible agents. However, the voluntary accrediting organizations (eg, AABB, FACT-JACIE) and some states or regions may require additional testing for allogeneic or autologous donors. See references for a list of selected publications containing testing requirements and standards. Required testing must be performed by a laboratory that is either certified to perform such testing on human specimens under the Clinical Laboratory Improvement Amendments of 1988 (42 USC §263a and 42 CFR 493) or has met equivalent requirements as determined by the Centers for Medicare and Medicaid Services.

[‡]Establishments not using FDA-licensed screening tests for HIV-1 group O antibodies must evaluate donors for risk associated with HIV-1 group O infection.

HPC(M) = hematopoietic progenitor cells from marrow; HPC(A) = HPCs from apheresis; MNC(A) = mononuclear cells from apheresis; NC(M) = nucleated cells from marrow; HPC(CB) = HPCs from cord blood; NC(CB) = NCs from cord blood; NC(WB) = NCs from whole blood; MS = maternal sample.

Table 1B. EU Minimal Requirements for Testing for Transmissible Agents in Cellular Therapy Products*^{†‡}

	Donors of HPC(M) and HPC(A)	Donors of HPC(CB), [§] NC(CB), MNC(A), NC(M), and NC(WB)
Timing of specimen collection	Up to 30 days before collection	Day of or up to 7 days after delivery
Human immunodeficiency virus, type 1 and type 2 (HIV-1, HIV-2)	X	X
Hepatitis B virus (HBV)	X	X
Hepatitis C virus (HCV)	X	X
<i>Treponema pallidum</i> (syphilis)	X	X
Human T-cell lymphotropic virus, type I (HTLV-I)	X	X

*The tests must be carried out by a qualified laboratory authorized as a testing center by the competent authority in the Member State, using CE-marked testing kits where appropriate. The type of test used must be validated for the purpose in accordance with current scientific knowledge. In certain circumstances, additional testing may be required depending on the donor's history and the characteristics of the tissue or cells donated (eg, malaria, toxoplasma, cytomegalovirus, Epstein-Barr virus, *Trypanosoma cruzi*).

[†]Member countries of the European Union may amend and/or introduce additional requirements. In some settings, testing by more than one method may be required for some infectious agents. This table is not intended to reflect all national variations but rather to present general requirements within the EU directives.

[‡]The governmental regulations require testing of autologous donors for transmissible agents only if removed cells are to be stored or cultured. However, the voluntary accrediting organizations (eg, AABB, FACT-JACIE) may require testing of autologous donors.

[§]The testing is repeated on the CBU if the CBU is stored for a long period of time; alternatively, nucleic acid testing (NAT) technology is used.

^{||}Performed on donors living in or originating from high-prevalence areas, or with sexual partners originating from those areas, or whose parents originate from those areas.

HPC(M) = hematopoietic progenitor cells from marrow; HPC(A) = HPCs from apheresis; MNC(A) = mononuclear cells from apheresis; NC(M) = nucleated cells from marrow; NC(WB) = NCs from whole blood; HPC(CB) = HPCs from cord blood; NC(CB) = NCs from cord blood; MS = maternal sample; CBU = cord blood unit.

See references for a list of selected publications containing testing requirements and standards.

*Application of biohazard and warning labels extends outside the HCT/Ps described in 21 CFR 1271 based on voluntary adherence to professional standards and applies to all products defined in this *Circular*, including HPC(M), which is not regulated under 21 CFR 1271. FDA eligibility processes and associated biohazard and warning labels were required on and after May 25, 2005 and may or may not be implemented for units collected before this date per facility-specific policy.

†When abnormal results of any donor screening or testing are identified in the donor, the transplant physician is notified of those results.

‡Urgent medical need must be documented when a donation is used for transplantation from a related or an unrelated donor with an “incomplete” eligibility status, or when a donation is used for transplantation from an unrelated donor with an “ineligible” eligibility status. When an HCT/P is made available for use before the donor eligibility determination is completed under the urgent medical need provision, the physician using the HCT/P must be notified that the testing and screening were not complete, and the notification must be documented [21 CFR 1271.60(d)(3)].

§Donor eligibility status of “incomplete” means donor eligibility determination was not completed per US requirements. The donor eligibility determination must be completed for donor screening and testing per FDA requirements during or after the use of the cellular therapy product. When not feasible to complete screening and/or testing per FDA criteria, documentation should be on file to justify. The biohazard label may be applied per facility-specific guidance for “incomplete” eligibility status.

¶FDA does not require donor testing or screening for autologous donors; FDA requires all autologous donations to be labeled “For Autologous Use Only.” When all donor screening and testing is not completed per FDA requirements, the label “Not Evaluated for Infectious Substances” is also required. Any abnormal donor screening or testing results (even though neither screening nor testing is mandated for this group of donors) require appropriate labeling [21 CFR 1271.90(a)(b)].

HCT/Ps = human cells, tissues, and cellular and tissue-based products; CFR = Code of Federal Regulations; HPC(M) = hematopoietic progenitor cells from marrow; FDA = Food and Drug Administration; NR = not required; R = required; NA = not applicable.

References

1. AABB, American Red Cross, America's Blood Centers, Armed Services Blood Program. Circular of information for the use of human blood and blood components. Bethesda, MD: AABB, 2013.
2. Food and Drug Administration. Current good tissue practice for human cell, tissue, and cellular and tissue-based product establishments; inspection and enforcement; final rule (November 24, 2004). Fed Regist 2004;69:68612-88.
3. Food and Drug Administration. Eligibility determination for donors of human cells, tissues, and cellular and tissue-based products; final rule (May 25, 2004). Fed Regist 2004;62:29786-834.
4. Food and Drug Administration. Human cells, tissues, and cellular and tissue-based products; donor screening and testing, and related labeling; interim final rule (May 25, 2005). Fed Regist 2005;70:29949-52.
5. Food and Drug Administration. Human cells, tissues, and cellular and tissue-based products; establishment registration and listing; final rule (January 19, 2001). Fed Regist 2001;66:5447-69.
6. Directive 2004/23/EC of the European Parliament and of the Council of 31 March 2004 on setting standards of quality and safety for the donation, procurement, testing, processing, preservation, storage and distribution of human tissues and cells. Official Journal of the European Union 2004;L102:48-58.
7. Commission Directive 2006/17/EC of 8 February 2006 implementing Directive 2004/23/EC of the European Parliament and of the Council as regards certain technical requirements for the donation, procurement and testing of human tissues and cells. Official Journal of the European Union 2006;L038:40-52.
8. Rice B, ed. Standard Terminology for Blood, Cellular Therapy, and Tissue Product Descriptions. Version 6.X. San Bernardino, CA: ICCBBA.
9. Distler P, ed. United States consensus guidance for the uniform labeling of cellular therapy products using ISBT 128. Version 1.3.1. San Bernardino, CA: ICCBBA, 2015. [Available at <http://www.iccbba.org/tech-library/iccbba-documents/guidance-documents> (accessed April 13, 2016).]
10. Distler P, ed. Implementation Guide: Product Coding [Data Structures 003 and 032] - Cellular Therapy, Version 1.2.0, San Bernardino, CA: ICCBBA, 2015.
11. Food and Drug Administration. Guidance for industry: Eligibility determination for donors of human cells, tissues, and cellular and tissue-based products (HCT/Ps). (August 8, 2007) Rockville, MD: CBER Office of Communications, Outreach, and Development, 2007.
12. Allickson J, ed. Standards for cellular therapy services. 7th ed. Bethesda, MD: AABB, 2015.
13. Standards for tissue banking. 13th ed. McLean, VA: American Association of Tissue Banks, 2012.
14. FACT-JACIE international standards for cellular therapy product collection, processing, and administration. 6th ed. Omaha, NE: Foundation for the Accreditation of Cellular Therapy, 2015.
15. NetCord-FACT international standards for cord blood collection, processing, and release for administration. 5th ed. Omaha, NE: Foundation for the Accreditation of Cellular Therapy, 2013.
16. NMDP standards. 23rd ed. Minneapolis, MN: National Marrow Donor Program, 2016.
17. Couriel D, Caldera H, Champlin R, Komanduri K. Acute graft-versus-host disease: Pathophysiology, clinical manifestations, and management. Cancer 2004;101:1936-46.

Specific Product Information

This page is intended to be blank to provide space for the distributing institution to provide additional product information as applicable to its product.

As indicated in the “General Information” section, the distributing institution is responsible for providing specific information not already included in this *Circular* about the cellular therapy product, including but not limited to the following:

- Description.
- Action.
- Indications.
- Contraindications.
- Storage.
- Dosage.
- Administration.

Note: Specific product labeling information is required if the product is manufactured under a US FDA-approved IND application or an IDE in the United States. Products manufactured and administered outside the United States must comply with local regulations.

This page is intended to be blank. The distributing facility can use this space to provide more specific information about the cellular therapy product.

This page is intended to be blank. The distributing facility can use this space to provide more specific information about the cellular therapy product.

This page is intended to be blank. The distributing facility can use this space to provide more specific information about the cellular therapy product.