

ADENOVIRUS INFECTION in Reptiles

ANIMAL GROUP AFFECTED	TRANSMISSION	CLINICAL SIGNS	FATAL DISEASE?	TREATMENT	PREVENTION & CONTROL
Lizards Snakes Also detected in chelonians and crocodiles	Most likely virus excreted in the faeces and ingested Respiratory droplets Vertically Inapparent carriers in adults	Unspecific Poor doers Poor appetite Sometimes diarrhea Sudden death Young lizards, 4 to 12 weeks old more affected	Not always. Inapparent carriers in adults	None against the virus Control of secondary bacterial infections, supportive treatment	Quarantine, PCR testing of newly introduced animals, hygiene.

Fact sheet compiled by Marja J. L. Kik, D.V.M., Ph.D, Diplomate vet Path, Reptile Veterinarian, Nieuwegein, The Netherlands	Last update April 2009
Fact sheet reviewed by Tibor Papp, Universität Hohenheim, Institut für Umwelt- und Tierhygiene, Stuttgart, Germany Rachel E. Marschang, Universität Hohenheim, Institut für Umwelt- und Tierhygiene, Stuttgart, Germany	
Susceptible animal groups: Most commonly found in squamata lizards. Agamid lizards: <i>Pogona vitticeps</i> (bearded dragon), <i>Pogona henrylawsoni</i> (Rankin's dragon), <i>Physignatus</i> spp. (water dragon). Chameleons: <i>Chamaelo jacksoni</i> (Jackson's chameleon), <i>Chamaelo montium</i> (mountain chameleon). Gekkota: <i>Hemitheconyx caudicinctus</i> (fat-tail gecko), <i>Eublepharus macularis</i> (leopard gecko), <i>Gekko gecko</i> (tokay gecko). Helodermatids: <i>Heloderma suspectum</i> (Gila monster), <i>Heloderma horridum</i> (Mexican beaded lizard). Monitors: <i>Varanus exanthematicus</i> (savannah monitor), <i>Varanus prasinus</i> (emerald monitor). Skinks: <i>Tiliqua scincoides intermedia</i> (blue-tongued skink), Also often detected in snakes. Boids: <i>Boa constrictor</i> (common boa), <i>Lichanura trivergata</i> (rosy boa). Viperids: <i>Crotalus scutulatus scutulatus</i> (Mojave rattlesnake), <i>Bothriechis marchi</i> (palm viper), <i>Vipera aspis aspis</i> , (asp viper). Colubrids: <i>Elaphe guttata</i> (corn snake), <i>Lampropeltis zonata multicincta</i> (Sierra mountain kingsnake), <i>Lampropeltis getula californiae</i> (California kingsnake), <i>Pituophis catenifer</i> (Pacific gopher snake). Recent reports from chelonians: <i>Inotestudo forsteni</i> (Sulawesi tortoise), <i>Trachemys scripta elegans</i> (red-eared slider), <i>Terrapene ornata ornata</i> (ornate box turtle) Earlier single report from crocodiles: <i>Crocodylus niloticus</i> (Nile crocodile)	
Causative organism Family <i>Adenoviridae</i> , mainly genus <i>Atadenovirus</i> , also genus <i>Siadenovirus</i> and unclassified (only in turtles)	
Zoonotic potential None	
Distribution World-wide	
Transmission Direct (fecal-oral), environmental (droplets), vertical (suspected to be through the egg in utero, or at time of oviposition)	
Incubation period Not known	
Clinical symptoms Unspecific, poor growth, poor appetite, sometimes diarrhea, sudden death, dermatitis in snakes. Mostly young animals, 4 to 12 weeks old, are affected. In snake adult animals can be affected too.	
Post mortem findings Possible findings: gastro-enteritis, stomatitis, oesophagitis, hepatitis, nephritis, pneumonia and encephalitis. Many bearded dragons don't show significant gross lesions.	



Microscopically: hepatic necrosis often with inflammatory cells in the sinusoids. Large basophilic intranuclear inclusion bodies in the hepatocytes and Kupffer cells. stomatitis, esophagitis, gastro-enteritis with intranuclear inclusion bodies in the epithelium. Intranuclear inclusion bodies in glial cells and endothelial cells in the brain.

Diagnosis

Polymerase Chain Reaction (PCR), in situ hybridization, histology (basophilic intranuclear inclusion bodies), transmission electron microscopy, virus isolation (if possible). Serological testing for antibodies against adenoviruses has been described in snakes using a neutralization test.

Material required for laboratory analysis

Fresh cloacal swabs, feces or tissues (liver best, also lung, kidney, intestine, etc.) at best in medium or saline solution containing antibiotics or frozen for PCR and virus isolation. Formalin fixed material (liver as well as other tissues, e.g. intestine) for histology and possibly also for PCR.

Relevant diagnostic laboratories

Rachel Marschang, DVM, PhD, FTÄ Microbiology, ZB. Reptilien, University Hohenheim, Institut für Umwelt und Tierhygiene, Garbenstr. 30, 70599 Stuttgart, Germany
Silvia Blahak, DVM, PhD, Chemisches und Veterinäruntersuchungsamt OWL, Westerfeldstrasse 1, 32758 Detmold, Germany
Balázs Harrach, DVM, PhD, DSc. Veterinary Medical Research Institute, Hungarian Academy of Sciences, H-1143 Budapest, Hungária krt. 21, Hungary
Szilvia Farkas, DVM, PhD, Veterinary Medical Research Institute, Hungarian Academy of Sciences, H-1143 Budapest, Hungária krt. 21, Hungary

Treatment

Use of antibiotics to control secondary infections. E.g. enrofloxacin, marbofloxacin, trimetoprim-sulfamethoxazole. Supportive treatment.

Prevention and control in zoos

- Quarantine new reptiles for a minimum of 90 days in a separate room, with separate set of husbandry tools, separate air duct system, use footbaths with virucidal disinfectant (e.g. bleach) at entrance. Weigh the animals as they enter and exit the quarantine. House the animals individually.
- Necropsy all animals that are euthanised or die.
- Check for internal and external parasites; treat the animals against these common parasites. Disinfect used materials and housing on a regular basis

Suggested disinfectant for housing facilities

Virucidal disinfectants, e.g. 10% chlorine bleach solution, quaternary ammonium compounds, ammonia.

Notification

Guarantees required under EU Legislation

Guarantees required by EAZA Zoos

Measures required under the Animal Disease Surveillance Plan

Measures required for introducing animals from non-approved sources

Measures to be taken in case of disease outbreak or positive laboratory findings

Conditions for restoring disease-free status after an outbreak

Contacts for further information

References

1. Benkő, M., Élő, P., Ursu, K., Ahne, W., LaPatra, S.E., Thomson D., Harrach B. 2002. First molecular evidence for the existence of distinct fish and snake adenoviruses. *J Virol.* 76 (19): 10056-9.
2. Benkő, M., Harrach, B., Both, G.W., Russell, W.C., Adair, B.M., Ádám, É., de Jong, J.C., Hess, M., Johnson, M., Kajon, A., Kidd, A.H., Lehmkuhl, H.D., Li, Q.G., Mautner, V., Pring-Akerblom, P., Wadell, G. 2005. Family Adenoviridae. In: Fauquet, C.M. (Ed.), *Virus Taxonomy. Eighth Report of the International Committee on Taxonomy of Viruses*, Academic Press, New York, pp. 213-228.
3. Farkas, S.L., Harrach, B., Benkő, M. 2008. Completion of the genome analysis of snake adenovirus type 1, a representative of the reptilian lineage within the novel genus *Atadenovirus*. *Vir. Res.* 132, 132-139.
4. Farkas, S.L., Gál, J. 2009. Adenovirus and mycoplasma infection in an ornate box turtle (*Terrapene*



- ornata ornata*) in Hungary. Vet. Microbiol. In press.
5. Heldstab A., Bestetti G. 1984. Virus associated gastrointestinal diseases in snakes. J Zoo Anim Med 15:118-28.
 6. Jacobson, E.R., Gardiner, C.H. 1990. Adeno-like virus in esophageal and tracheal mucosa of a Jackson's chameleon (*Chamaeleo jacksoni*). Vet. Path. 27: 210-212.
 7. Jacobson, E.R., Gaskin, J.M., Gardiner, C.H. 1985. Adenovirus-like Infection in a Boa Constrictor J.A.V.M.A. 187:1226-7.
 8. Jacobson, E.R., Gardiner, C.H., Foggin, C.M. 1984. Adenovirus-like infection in two Nile crocodiles. J Am Vet Med Assoc. 185: 1421-22.
 9. Jacobson, E.R., Kollias, G.V. 1986. Adenovirus-like infection in a savannah monitor. J. Zoo Anim. Med. 17: 149-151.
 10. Jacobson, E.R., Kopit, W, Kennedy, F.A., Funk, R.S. 1996. Coinfection of a bearded dragon, *Pogona vitticeps*, with adeno- en dependovirus-like viruses. Vet Pathol 33:343-346.
 11. Kim, D.Y., L.E., Mitchell, M.A., Bauer, R.W., Poston, R. Cho, D.Y. 2002. An outbreak of adenoviral infection in inland bearded dragon (*Pogona vitticeps*) coinfectd with dependovirus and coccidial protozoa (*Isospora sp.*). J Vet Diagn Invest. 14 (4): 332-4.
 12. Marschang, R.E., Mischling, M., Benkő, M., Papp, T., Harrach, B., Böhm, R. 2003. Evidence for wide-spread atadenovirus infection among snakes. In: Jestin, A., Clement, G. (Eds.), Virus persistence and evolution. Proceedings of the 6th International Congress of Veterinary Virology. ZOO-POLE développement –ISPAIA, Ploufragan, France. p. 152.
 13. Ogawa, M. Ahne, W. Essbauer S. 1992. Reptilian viruses: adenovirus-like agent isolated from royal python (*Python regius*). Zentralbl. Verterinarmed. B. 39 (10): 732-736.
 14. Papp, T., Fledelius, B., Schmidt, V., Kaján, G.L., Marschang, R.E. 2009. PCR-sequence characterization of new adenoviruses found in reptiles and the first successful isolation of a lizard adenovirus. Vet. Microbiol. 134: 233-240.
 15. Perkins, L.E., Campagnoli, R.P., Harmon, B.G., Gregory C.R., Steffens W.L., Latimer, K., Clubb, S., Crane, M. 2001. Detection and confirmation of reptilian adenovirus infection by in situ hybridization. J Vet Diagn Invest. 13 (4): 365-8.
 16. Ramis, A., Fernandez-Bellon, H., Majo, N., Martinez-Silvestre, A., Latimer, K., Campagnoli, R., Harmon, B.G., Gregory, C.R., Steffens, W.L., Clubb, S., Crane, M. 2000. Adenovirus hepatitis in a boa constrictor (*Boa constrictor*). J. Vet. Diagn. Invest. 12: 573-576.
 17. Raymond, J.T., Lamm, M., Nordhausen, R. Latimer, K., Garner, M.M. 2003. Degenerative encephalopathy in a coastal mountain kingsnake (*Lampropeltis zonata multifasciata*) due to adenoviral-like infection. J. Wildl. Med. 39 (2): 431-436.
 18. Rivera, S., Wellehan, J.F.X., McManamon, R., Innis, C.J., Garner, M.M., Raphael, B.L., Gregory, C.R., Latimer, K.S., Rodriguez, C.E., Diaz-Figueroa, O., Marlar, A.B., Nyaoke, A., Gates, A.E., Gilbert, K., Childress, A.L., Risatti, G.R., Frasca S. 2008. Systemic adenovirus infection in Sulawesi Tortoises (*Indotestudo fortseni*) caused by a novel siadenovirus. 15th annual conference of Association of Reptilian and Amphibian Veterinarians. Los Angeles, CA, USA, October 12-15.
 19. Schumacher, J., Jacobson, E.R., Burns, R., Tramontin, R.R. 1994. Adenovirus-like infection in two rosy boas (*Lichanura trivirgata*). J. Zoo and Wildlife Med. 25: 461-465.
 20. Wellehan, J.F.X., Johnson, A.J., Harrach, B., Benkő, M., Pessier, A.P., Johnson, C.M., Garner, M.M., Childress, A., Jacobson, E.R. 2004. Detection and analysis of six lizard adenoviruses by consensus primer PCR provides further evidence of a reptilian origin for the atadenoviruses. J Virol. 78: 13366-9.
 21. Wozniak E.J., Denardo, D.F., Brewer, A., Wong, V, Tarara, R.P. 2000. Identification of adenovirus- and dependovirus-like agents in an outbreak of fatal gastroenteritis in captive born California mountain kingsnakes, *Lampropeltis zonata multicincta*. J Herpetol Med Surg 10 (2): 4-7.