

COLUMBIA-SUICIDE SEVERITY RATING SCALE

Screening Version - Recent

	Past Month	
	YES	NO
Ask questions that are bold and underlined		
Ask Questions 1 and 2		
1) Wish to be Dead: Person endorses thoughts about a wish to be dead or not alive anymore, or wish to fall asleep and not wake up. <u>Have you wished you were dead or wished you could go to sleep and not wake up?</u>		
2) Suicidal Thoughts: General non-specific thoughts of wanting to end one's life/die by suicide, <i>"I've thought about killing myself" without general thoughts of ways to kill oneself/associated methods, intent, or plan.</i> <u>Have you actually had any thoughts of killing yourself?</u>		
If YES to 2, ask questions 3, 4, 5, and 6. If NO to 2, go directly to question 6		
3) Suicidal Thoughts with Method (without Specific Plan or Intent to Act): Person endorses thoughts of suicide and has thought of a least one method during the assessment period. This is different than a specific plan with time, place or method details worked out. <i>"I thought about taking an overdose but I never made a specific plan as to when where or how I would actually do it...and I would never go through with it."</i> <u>Have you been thinking about how you might kill yourself?</u>		
4) Suicidal Intent (without Specific Plan): Active suicidal thoughts of killing oneself and patient reports having <u>some intent to act on such thoughts</u> , as opposed to <i>"I have the thoughts but I definitely will not do anything about them."</i> <u>Have you had these thoughts and had some intention of acting on them?</u>		
5) Suicide Intent with Specific Plan: Thoughts of killing oneself with details of plan fully or partially worked out and person has some intent to carry it out. <u>Have you started to work out or worked out the details of how to kill yourself and do you intend to carry out this plan?</u>		

	Past 3 Months	
	YES	NO
6) Suicide Behavior <u>Have you done anything, started to do anything, or prepared to do anything to end your life?</u> Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump; or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc.		

For inquiries and training requirements contact: Kelly Posner, Ph.D.

New York State Psychiatric Institute, 1051 Riverside Drive, New York, New York, 10032; posnerk@nyspi.columbia.edu

© 2008 The Research Foundation for Mental Hygiene, Inc.